



Relationship between maxillary disjunction and level of asthma control in school-aged patients

Relación entre disyunción maxilar y nivel de control del asma en pacientes de edad escolar

Gilberto Sáenz-Guerrero,* Rosaura Pacheco-Santiesteban,* Mario Soto-Ramos,[‡]
Humberto Alejandro Monreal-Romero,* Guillermo Martínez-Mata*

*Universidad Autónoma de Chihuahua. Chihuahua, Mexico.

[‡]Hospital Infantil de Especialidades de Chihuahua, Hospital Ángeles Chihuahua. Mexico.

ABSTRACT. Introduction: Dentofacial disorders in children are associated significantly to respiratory disorders, particularly asthma and allergic rhinitis, and although not pinpointed the mechanism by which impact these pathologies craniofacial development, it is suggested that obstruction upper airway difficult nasal breathing forcing the patient to compensate such obstruction by mouth breathing, this condition being the origin of malocclusion. **Objective:** The present study was conducted to determine whether treatment of malocclusion, specifically the correction of transverse compression maxillary positively influences asthma control assessed by clinical evidence of asthma control (Asthma Control Test) and as spirometry (FEV₁, FEV₁/FVC) at baseline and response tests bronchodilator addition of drug treatment. **Material and methods:** The study just took a group of 15 patients between seven and 12 years of age diagnosed with asthma and allergic rhinitis which were under pharmacological medical treatment for a minimum of six months. **Results:** Among the main results it was observed that there is a significant correlation between the maxillary disjunction and clinical asthma control and lung function in children with asthma and allergic rhinitis, mainly from the third month of orthopedic treatment. **Conclusion:** The use of maxillary circuit breakers, represents an alternative for an adequate management of respiratory disorders manipulating the conditions in the generation processes making the proposed system more efficient.

Keywords: Asthma, rhinitis, maxillary disjunction.

RESUMEN. Introducción: Las alteraciones dentofaciales en la edad pediátrica se encuentran asociadas de manera importante a trastornos respiratorios, principalmente al asma y la rinitis alérgica, y aunque no está claramente establecido el mecanismo por el cual impactan dichas patologías del desarrollo craneofacial, se sugiere que la obstrucción de la vía aérea superior dificulta la respiración nasal obligando al paciente a compensar dicha obstrucción por medio de una respiración bucal, siendo esta última el origen de la maloclusión. **Objetivo:** El presente estudio tuvo como objetivo determinar si el tratamiento de la maloclusión, específicamente la corrección de la compresión transversal del maxilar superior, influye positivamente en el control del asma evaluado por la prueba clínica de control del asma (Asthma Control Test), así como por espirometría (FEV₁, FEV₁/FVC) en pruebas basales y respuesta a broncodilatador, además del tratamiento farmacológico. **Material y métodos:** El estudio se llevó a cabo en un grupo de 15 pacientes entre los siete y 12 años de edad con diagnóstico de asma y rinitis alérgica, los cuales se encontraban bajo tratamiento médico farmacológico por un tiempo mínimo de seis meses. **Resultados:** Se observó una correlación significativa entre la disyunción maxilar y el control clínico del asma y la rinitis alérgica, mejorando de manera importante la función pulmonar a partir del tercer mes del tratamiento de ortopedia dentomaxilar. **Conclusión:** El uso de disyuntores maxilares representa una alternativa para un adecuado manejo de desórdenes respiratorios, manipulando las condiciones en los procesos de generación haciendo el sistema propuesto más eficiente.

Palabras clave: Asma, rinitis, disyunción maxilar.

Correspondence:

Humberto Alejandro Monreal-Romero, MD

Universidad Autónoma de Chihuahua. Chihuahua, Mexico.

E-mail: hmonreal@uach.mx

Received: VII-01-2021; accepted: III-24-2022.

How to cite: Sáenz-Guerrero G, Pacheco-Santiesteban R, Soto-Ramos M, Monreal-Romero HA, Martínez-Mata G. Relationship between maxillary disjunction and level of asthma control in school-aged patients. Neumol Cir Torax. 2022; 81 (1): 19-25.

INTRODUCTION

In order to understand the close relationship between respiratory function and the development of the stomatognathic apparatus, we must remember that the function is closely related to growth and development, in such a way that through nasal breathing multiple nerve endings are stimulated that generate diverse responses such as the amplitude of thoracic movement, three-dimensional development of the nostrils, ventilation and the size of

the maxillary sinuses, but which also induce the growth and remodeling of the adjacent orofacial structures, as well as the transverse development of the maxilla and the direction of facial growth.¹ Some researchers claim that nasal obstruction is associated with chronic mouth breathing, constituting the main etiological factor of anomalous craniofacial development, manifesting clinically as vertical facial growth or long face syndrome with subsequent dentomaxillary deformities.² Among the main stomatological manifestations, deep and narrow palates characterized by inverted V-shaped arches and an increase in palatal depth are reported, with a consequent maxillary protrusion, developing in most cases a class II subdivision 2 malocclusion, also known as distal dysgnathia, associated with mandibular retrusion.^{3,4} In addition, mechanical obstructions of congenital origin have been found, such as choanal atresia, alterations in Waldeyer's ring and other endonasal obstructions, as well as physiological alterations of the airways, which if not resolved in time may cause dentocraniofacial deformities.⁴ The prevalence of asthmatic problems has been increasing during the last years, rising from 3.2 to 25%, being considered the most frequent chronic respiratory disease.⁵ In Mexico, asthma affects from 5 to 8% of the population, with variations among the different regions of the country.⁶ The prevalence of asthmatic problems has been on the rise in recent years, increasing from 3.2 to 25%, being considered the most frequent chronic respiratory disease.⁵ In Mexico, asthma affects 5 to 8% of the population, with variations among the different regions of the country.⁶ Thus, the states of Colima, Tabasco, Chihuahua, Yucatán and Mexico City are those with the highest incidence figures of bronchial asthma.^{7,8} Multiple epidemiological, pathophysiological and therapeutic studies have demonstrated the association between allergic rhinitis and asthma; 90% of patients diagnosed with asthma have allergic rhinitis,⁹ making it the most frequent chronic disease¹⁰ and the most common childhood allergic disease.¹¹ It is clinically defined as a symptomatic disorder of the nose induced by immunoglobulin E (IgE)-mediated inflammation of the membranes lining the nose following exposure to an allergen.¹⁰ Asthma and allergic rhinitis are chronic respiratory diseases that obstruct the upper airways,¹² inducing an alternative mode of breathing and contributing to breathing through the mouth,⁷⁻⁹ resulting in orofacial developmental disorders, with dental malocclusion being the most frequent manifestation and of greatest interest to the pediatric stomatologist.¹³ The development of the craniofacial mass and specifically the components of the stomatognathic apparatus may be altered in the presence of a nasorespiratory obstruction, since if nasal breathing is compromised, it is compensated by oral breathing,¹⁴ which causes an imbalance between the lips, buccinator

muscles and tongue, exerting an abnormal force on the vestibular and lingual faces of the teeth, resulting in dental malpositions, mainly in the transversal direction.¹⁵

However, the severity of asthma can change over time and depends not only on the severity of the underlying disease but also on its response to treatment.¹⁶ Asthma and allergic rhinitis have been defined as chronic inflammatory airway diseases, which are closely related, especially in childhood, and are manifested by both upper and lower airway obstruction, emphasizing that for good asthma control it is important to adequately control allergic rhinitis, as lack of control of the latter leads to a suboptimal level of asthma control.¹⁷ In the most recent revision of the global initiative for the treatment of asthma GINA 2021, it is established that asthma should be evaluated periodically to verify the level of control of the disease and it is classified as: completely controlled, partially controlled or not controlled at all, as well as to perform pulmonary function measurements by means of spirometry or flowmetry, which allows determining the risk of possible later complications. Likewise, in cases where good control is not achieved, it is recommended to investigate adherence to treatment and the correct technique for the use of inhaled medications, as well as adverse environmental factors, comorbidities such as allergic rhinitis, rhinosinusitis, obesity, among others.

The objective of this research work was to establish how pediatric patients with a diagnosis of asthma and with strict pharmacological control of asthma, who present long face syndrome due to being chronic mouth breathers, can improve their pulmonary function by receiving a dentofacial orthopedic treatment, specifically, rapid maxillary expansion.

MATERIAL AND METHODS

The sample consisted of 15 patients who attended the Pediatric Pneumology Service of the Hospital Infantil de Especialidades del Estado de Chihuahua, aged between 7 and 12 years, all of whom had a baseline diagnosis of asthma and allergic rhinitis. These patients were under strict pharmacological control; However, when they were checked by the pediatric stomatologist, long face syndrome and maxillary compression in a transversal sense were observed, which we consider to be a factor that can interfere in the maximum results that can be obtained with the pharmacological medical treatment, for which reason, with previous authorization of the parents, each one of them was given a stomatological file that included diagnostic aids such as panoramic X-rays and study models, as part of the treatment and follow-up by the Pneumology Service, the patients were evaluated on four occasions, with a time interval of 30 days between one evaluation and another. The first intervention consisted in the evaluation




Figure 1:

A) A Hyrax® type rapid maxillary expansion screw is shown. **B)** It can be seen how the transverse width of the maxilla is calculated with the intention of obtaining the dentomaxillary discrepancy.

of asthma control by the treating pulmonologist, applying the Asthma Control Test (ACT), with the presence of the pediatric stomatologist to determine if despite being well controlled the patient presented long face syndrome and transverse compression of the maxilla.

The following protocol was followed during the four visits: first the ACT test was performed, then spirometry was performed with an EasyOne® ultrasonic spirometer to determine the values of forced vital capacity (FVC), forced expiratory volume in the first second (FEV1), as well as the FEV1/FVC ratio, and finally a bronchodilator was applied, and after 20 minutes the spirometry was repeated in a comparative manner.

- Pharmacological treatment. All patients who participated in the study were under the following pharmacological scheme for asthma: an inhaled steroid was used in combination with a long-acting β_2 agonist (salmeterol/fluticasone) in metered-dose aerosol, at a low dose of fluticasone, that is, a bottle containing a concentration of 25/50 μg , to receive in total 100/200, the total dose being 50/100 μg every 12 hours of salmeterol/fluticasone. As for the treatment of allergic rhinitis, this consisted of the application of a nasal spray of mometasone fuorate 0.05%, with 50 μg per dose every 24 hours in both nostrils.
- Dentomaxillary orthopedic treatment. The orthopedic treatment consisted of placing an 11 mm Hyrax® type fixed palatal disjunct, adjusting universal steel Dentaurem® orthodontic bands in upper first permanent molars and impressions were taken with Kromopan® alginate, impressions were taken with Kromopan® alginate, which were cut in high resistance stone plaster type III Magnum®, thus obtaining a working model, on

which the fast maxillary disjunction appliance was made, to finally be cemented in the patient's oral cavity with a reinforced glass ionomer AquaCem®. The amount of activation was indicated to each patient according to the degree of maxillary compression estimated by means of the space analysis proposed by Pont, which consists of estimating the arch perimeter necessary for dental eruption based on the mesiodistal width of the upper permanent incisors and upper permanent lateral incisors (Figure 1). Once the expansion was completed, the appliance was left for three months without activation, acting as a retainer, thus allowing the ossification of the palatal suture at the site of disjunction, reducing the possibility of recurrences. Once the ossification of this area was corroborated by taking an occlusal radiograph, the appliance was removed. The data obtained were analyzed using SPSS version 2021 statistical software. The present work was carried out in accordance with the codes of ethics of the world medical association for experiments, including human experiments.

RESULTS

ACT test. Of the patients who attended the pulmonology office of the Hospital Infantil de Especialidades de Chihuahua, 15 participated in this research protocol, who were under strict pharmacological control, with a minimum treatment time of six months, such treatment was focused on the control of asthma and allergic rhinitis, the sample consisted of 10 male patients and five female patients, between the age ranges of seven to 12 years, with a mean of 8.46 and a SD of 1.50 years. The age at which patients were diagnosed as asthmatic ranged from five to 96 months, with a mean age at diagnosis of

39.53 months (3.2 years) and a SD of 24.60 months (2.05 years). All patients underwent a space analysis, obtaining the degree of transverse maxillary compression, where a mean compression in the anterior segment of 4.6 mm was observed, and in the posterior segment of 5.2 mm, with a SD of 1.5 and 2.4 mm, respectively. It is important to mention that when there is a transverse compression of the maxilla, the palate becomes deeper at the expense of the floor of the nostrils. An anterior maxillary compression standard deviation of 1.5 mm and a posterior maxillary compression standard deviation of 2.4 mm was obtained.

Based on the results obtained in the ACT test, it was observed that all the patients in the sample had good or acceptable control of their clinical condition, since their score was equal to or greater than 20 points as an effect of the continuous pharmacological treatment they had at least six months before entering the study. However, the patients clinically manifested adenoid facies and long face syndrome, a sign that the patient is a mouth breather, which almost always causes maxillary compression, a manifestation corroborated by the pediatric stomatologist

at the first visit, as previously mentioned. Spirometry studies were also performed in all visits to each of the cases as part of the objective evaluation of asthma control by measuring pulmonary function according to age, weight and height and thus FEV₁ and its predicted percentage, FVC and its predicted percentage and the FEV₁/FVC ratio were obtained. It should be noted that spirometry provides an objective measurement to determine airway obstruction in patients with asthma and to measure the level of severity of airway obstruction. The FEV₁/FVC ratio allows determining the presence of airway obstruction, since normally this index should be greater than 0.8. When it is less than 0.8, an obstructive pattern is defined in the spirometry and to classify the degree of obstruction the FEV₁ result is used compared to its predicted value, leaving this classification of obstruction as described in [Table 1](#).

The FEV₁ values obtained were equal to or greater than 85% of the predicted value from the first visit or visit zero ([Table 2](#)), indicating good asthma control due to strict adherence to pharmacological treatment; however, it can be observed that at visit three there was a significant increase in the same reaching values of 89.6% of the predicted FEV₁, here the variable included was the rapid decompression of the maxilla in a transverse direction. This allows us to suggest that such decompression favors pulmonary function. The above data were verified with a Pearson correlation where we found that the highest correlation was between the percentage predicted at visit two and visit three. There is a statistically significant increase from visit one, increasing and lasting the effect until visit three ([Table 3](#)).

As part of the protocol, spirometry with application of a bronchodilator was performed 20 minutes after the baseline

Table 1: Classification of the degree of obstruction obtained by spirometry.

FEV ₁ % of predicted value	Obstruction
70-80	Mild
60-69	Moderate
50-59	Moderately severe
35-49	Severe
< 35	Very severe

Table 2: Predicted percentage values for baseline spirometry at each visit for all patients.

FEV ₁ % predicted	Media ± standard deviation	Minimum	Maximum
Visit 0	86.108 ± 15.92256	63.79	110.89
Visit 1	85.97467 ± 14.07274	62.76	110.89
Visit 2	85.85067 ± 17.9075	51.44	111.83
Visit 3	89.608 ± 14.53569	65.33	113.10

Table 3: Correlation between variables with post BD FEV₁ and baseline FEV₁ values (Pearson correlation).

FEV ₁	%PRED/0	%PRED/1	%PRED/2	%PRED/3
%PRED/0	1.0000			
%PRED/1	0.6983	1.0000		
%PRED/2	0.7791	0.9096	1.0000	
%PRED/3	0.7096	0.8445	0.9163	1.0000

spirometry, for comparative purposes, by obtaining the average percentage change using the following formula:

$$(\text{FEV}_1 \text{ post BD} - \text{FEV}_1 \text{ baseline})$$

$$\text{Baseline FEV}_1 = \% \text{ change}$$

When the percentage change is greater than 12% there is a significant change and thus the volume measured in milliliters was using the following formula: $\text{FEV}_1 \text{ post BD} - \text{FEV}_1 \text{ basal}$, when it is more than 150 mL (200 mL), we have a significant change.

The mean percentage change was less than 12%, which indicates that there is no significant change after the application of the bronchodilator, which means that

most patients were well controlled pharmacologically, although between visits zero and one some cases showed an increase of more than 12%, which indicates that despite the pharmacological treatment at the time of maxillary expansion, an additional improvement in pulmonary function was obtained, as shown in cases 6 and 14 of [Table 4](#).

A Pearson correlation was made between the percentages of change and maxillary compression and the following was found: there is a statistically significant and positive correlation between anterior and posterior compression (0.77), which means that as the anterior maxillary compression increases, so does the posterior maxillary compression. The percentage of change between visit zero and visit one is also positive (as the percentage of change between visit zero increases in this sample so does that of visit one (0.70), that is, as maxillary expansion increases the percentage of change, this effect was also observed in visit two (0.59) although to a lesser degree and visit three (0.67) ([Table 5](#)).

Due to the great demographic variability in the country on the prevalence of asthma and rhinitis in children and adolescents, the results shown in this research are conclusive to represent a statistical behavior with positive bias, since it is more common to find values above the value of zero and not below zero, which would show the total absence of such clinical manifestations.

DISCUSSION

In the present study, the ratio of asthmatic patients corresponds to a ratio of 2:1 with respect to gender, that is, it is more frequent in the male gender in pediatric age. Most orthopedic treatments in asthmatic patients are aimed at increasing nasal airflow by disjunction of the maxilla, decreasing airflow resistance from 45 to 53%.¹⁷ When the mid-palatal suture is expanded, nasal airflow capacity is increased when measured at maximal effort.¹⁸ Other authors have evaluated the changes in nasal airflow after rapid maxillary expansion, with a significant increase in airflow through the nasal passage.¹⁹ It is important to mention that

Table 4: Average percent change per patient post BD.

Patient	Visit 1	Visit 2	Visit 3	Visit 4
1	5.83	1.77	1.76	3.42
2	0.79	1.43	6.95	1.43
3	0.47	7.92	11.48	4.43
4	2.76	8.19	1.78	0.56
5	6.17	4.50	2.48	6.43
6	8.13	11.85	2.96	19.85
7	11.66	11.40	4.54	5.09
8	5.9	7.93	8.71	7.93
9	1.27	1.29	2.56	1.94
10	4.25	8.69	4.66	6.21
11	3.91	3.16	1.25	5.30
12	3.75	17.70	4.70	8.69
13	3.31	2.44	3.33	2.44
14	26.05	21.81	15.05	13.97
15	3.40	0.33	5.26	2.30

Table 5: Pearson's correlation between percent change and maxillary compression post BD.

		Compression		% of change			
		Anterior	Posterior	Visita 0	Visita 1	Visita 2	Visita 3
Compression	Anterior	1.000					
	Posterior	0.7788	1.000				
% of change	Visit 0	-0.0996	-0.2330	1.000			
	Visit 1	0.2792	0.1928	0.7029	1.00		
	Visit 2	-0.1160	-0.2868	0.5308	0.5327	1.000	
	Visit 3	0.2049	-0.1017	0.5915	0.6795	0.2897	1.000

none of the studies found in the literature review performed a measurement like this, being that spirometry is a useful, reliable and accessible tool in tertiary hospitals such as the Hospital Infantil de Especialidades de Chihuahua. It is worth mentioning that the maximum improvement in pulmonary function measured by spirometry was observed three months after maxillary expansion, which had not been previously reported by other authors. In this study it is observed that the implementation of the maxillary expansion treatment significantly improved the level of clinical control of asthma as measured by the ACT questionnaire. To show the involvement of the disjunctur during expansion, we propose the following mechanism: due to the fact that disjuncture allows bone separation and this in turn exerts a tension on the palatal suture, a stimulation of osteoblastic activity is generated for bone formation function through the generation of mechanical stress between the sections of the maxillary bone with the consequent production of mediators such as prostaglandins, nitric oxide and growth factors that are able to overcome the resorption process.

CONCLUSIONS

It is essential to perform a multidimensional evaluation of the patient's asthma control, since the ACT test alone should not be considered reliable because it is a subjective test, since it depends on the patient's responses; however, when correlated with pulmonary function it acquires a greater value, since it is expected to have a normal ACT test with a good pulmonary function, the latter measured with baseline spirometry. Now, in this study, maxillary expansion was added as a therapeutic strategy with the intention of improving nasal breathing, annulling mouth breathing, so that once the expansion was performed, it could be observed that there was no longer a positive bronchodilator response, which was observed at visit zero, suggesting that prior to the expansion the patient could still improve his lung function, which at visits two and three approached optimal values.

According to the results obtained, maxillary expansion had a negative correlation with FEV1, that is, the greater the maxillary compression, the lower the FEV1 value. These data are important because they show that patients with greater upper airway obstruction have decreased lung function, as measured by spirometry.

REFERENCES

1. Taboada-Aranza O, Torres-Zurita A, Cazares-Martinez CE, Orozco-Cuanalo L. Prevalence of malocclusions and speech disorders in a preschool population in eastern Mexico City. *Bol Med Hosp Infant Mex*. 2011;68(6):425-430. Available in: www.scielo.org.mx/scielo.php?script=sci_abstract&pid=S1665
2. Lina-Moreno M, Sara-Howe C, Colleen-Kummet, Kaci-Vela C, Deborah-Dawson V, Thomas-Southard E. Phenotypic diversity in white adults with moderate to severe class II malocclusion. *AJO-DO*. 2014;145(3):305-316. doi: 10.1016/j.ajodo.2013.11.013.
3. Morales-Ugalde F. Classification of malocclusion in the anteroposterior vertical and transverse planes. *Rev ADM*. 2007;13(3):97-109. Available in: www.medigraphic.com/adm
4. Parra-Iraola S, Zambrano-Mendoza AG. Mouth deforming habits present in preschoolers and school children: systematic review. *Int J Odontostomat*. 2018;12(2):188-193. Available in: <http://dx.doi.org/10.4067/S0718-381X2018000200188>
5. Natera-Marcote A, Gasca-Vargas V, Rodriguez-Yañez E, Casasa-Araujo R. Treatment of deep bite-clinical case. *Rev Lat Ort*. 2006;2:1-10. Available in: <https://www.ortodoncia.ws/publicaciones/2005/art-16>
6. Gisenia-Iniquilla A, Tania-Padilla C, Sonia-Macedo C, Nedy-Hilari O. Relationship of dental malocclusion with body position and plant footprint in a group of aymaras teenagers. *Rev investig Altoandin*. 2017;19(3):255-264. Available in: <http://dx.doi.org/10.18271/ria.2017.290>
7. Dahan Ch, Boul-Serhal DDS, Englebert LSD. Cephalometric changes in class II, division 1 cases after orthopedic treatment with the bioactivator. *AJO-DO*. 1989;95(2):127-137. Available in: [https://doi.org/10.1016/0889-5406\(89\)90391-0](https://doi.org/10.1016/0889-5406(89)90391-0)
8. Giuseppe-Perinetti, Luca-Contardo, Silvestrini-Biavati A, Perdoni L, Attilio C. Dental malocclusion and body posture in young subjects: a multiple regression study clinics. *Clinics*. 2010;65(7):689-695. doi: 10.1590/S1807-59322010000700007.
9. Lucía-Bernal V, Harold M, Herrera-Yudi C, Montoya C. Characteristics dento skeletal with relation to corporal posture in class II malocclusion craniocervical posture in children with class I, II and III skeletal relationships. *Pesq Bras Odontoped Clin Integr*. 2017;17:1-12. doi:10.4034/PBOCI.2017.171.07.
10. Michelotti A, Buonocore G, Manzo P, Pellegrino G, Farella M. Dental occlusion and posture: an overview. *Prog orthod*. 2011;12(1):53-58. Available in: <https://doi.org/10.1016/j.pio.2010.09.010>
11. Lippold C, Danesh G, Schilgen M, Drerup B, Hackenberg L. Relationship between thoracic, lordotic, and pelvic inclination and craniofacial morphology in adults. *Angle Orthod*. 2006;76(5):779-785. doi: 10.1043/0003-3219(2006)076[0779:RBTLAP]2.0.CO;2.
12. Valentino B, Melito F. Functional relationship between the muscles of mastication and the muscles of the leg: an electromyographic study. *Surg Radiol Anat*. 1991;13:33-37. doi: 10.1007/BF01623138.
13. Attilio-Michelle D, Filippi-Maria R, Femminella B, Festa F, Tecco S. The influence of an experimentally-induced malocclusion on vertebral alignment in rats: a controlled pilot study. *Cranio*. 2005;23(2):119-129. doi: 10.1179/crn.2005.017.
14. Huggare J. Postural disorders and dentofacial morphology. *Acta Odontol Scand*. 1998;56(6):383-386. doi: 10.1080/0001635-98428374.
15. Festa F, Tecco S, Dolci M, Ciufolo F, Di-Meo S, Filippi-Maria R. Relationship between cervical lordosis and facial morphology in caucasian women with a skeletal class II malocclusion: a cross-sectional study. *Cranio*. 2003;21(2):121-129. doi: 10.1080/08869634.2003.11746240.
16. Marchena-Rodriguez A, Moreno-Morales N, Ramirez-Parga E, Labajo-Manzanarez M, Luque-Suarez A, Guijon-Nogueron G. Relationship between foot posture and dental malocclusions in children aged 6 to 9 years: a cross-sectional study. *Medicine*. 2018;97(19):1-9. doi: 10.1097/md.00000000000010701.

17. Soto-Ramos M, Gochicoa-Rangel L, Hinojos-Gallardo L, Hernández-Saldaña R, Baquera-Arteaga M. Evaluación multidimensional del control en el paciente con asma. Aplicaciones clínicas en la toma de decisiones para el tratamiento. Neumol Cir Torax. 2013;72(1):52-61. Available in: <https://www.medigraphic.com/cgi-bin/new/resumen.cgi?>
18. Fusun Y, Faruk A. Evaluating mandibular cortical index quantitatively. Eur J Dent. 2008;2(4):283-290. doi:10.1055/s-0039-1697393.
19. Sonnesen L, Kjaer I. Anomalies of the cervical vertebrae in patients with skeletal class II malocclusion and horizontal maxillary overjet. AJO-DO. 2008;133(2):15-20. doi: 10.1016/j.ajodo.2007.07.018.

Conflict of interests: The authors declare that they have no conflict of interests.